

EXOTIC SMOOTH STRUCTURES ON $3\mathbb{C}P^2\#8\overline{\mathbb{C}P}^2$

JONGIL PARK*

ABSTRACT. Motivated by Stipsicz and Szabó's exotic 4-manifolds with $b_2^+ = 3$ and $b_2^- = 8$, we construct a family of simply connected smooth 4-manifolds with $b_2^+ = 3$ and $b_2^- = 8$. As a corollary, we conclude that the topological 4-manifold $3\mathbb{C}P^2\#8\overline{\mathbb{C}P}^2$ admits infinitely many distinct smooth structures.

1. INTRODUCTION

Since S. Donaldson introduced gauge theory in 1982, it has been known that most simply connected smooth 4-manifolds with b_2^+ odd and large enough admit infinitely many distinct smooth structures ([4], [7], [15], [16]). But it is still an intriguing problem to determine which smooth 4-manifolds with b_2^+ small have more than one smooth structure. In the case when $b_2^+ = 1$, S. Donaldson first proved that a Dolgachev surface is not diffeomorphic to $\mathbb{C}P^2\#9\overline{\mathbb{C}P}^2$ ([2]) and D. Kotschick proved in the late 1980's that the Barlow surface is not diffeomorphic to $\mathbb{C}P^2\#8\overline{\mathbb{C}P}^2$ ([9]). Recently, the author constructed a new simply connected symplectic 4-manifold with $b_2^+ = 1$ and $b_2^- = 7$ ([17]), and then R. Fintushel, R. Stern, A. Stipsicz and Z. Szabó found many new exotic smooth 4-manifolds with $b_2^+ = 1$ using a rational blow-down technique ([5], [18], [21]). So it has been proved up to now that rational surfaces $\mathbb{C}P^2\#n\overline{\mathbb{C}P}^2$ with $n \geq 5$ admit infinitely many distinct smooth structures. On the other hand, the case with $b_2^+ = 3$ has also been studied extensively for a long time. For example, it was known in the mid 1990's that the $K3$ surface $E(2)$ and $3\mathbb{C}P^2\#19\overline{\mathbb{C}P}^2$ admit infinitely many distinct smooth structures ([3], [6], [10], [20]) and the topological 4-manifolds $3\mathbb{C}P^2\#n\overline{\mathbb{C}P}^2$ with $n \geq 14$ also admit infinitely many distinct smooth structures ([19], [23]). And later, the same statement with $n \geq 10$ was also proved ([11], [12], [13]). Recently, Stipsicz and Szabó constructed infinitely many distinct smooth structures on $3\mathbb{C}P^2\#9\overline{\mathbb{C}P}^2$ and they also constructed infinitely many exotic smooth 4-manifolds with vanishing first homology, $b_2^+ = 3$ and $b_2^- = 8$ using a similar technique appeared in [18] ([22]).

The aim of this paper is to prove that the topological 4-manifold $3\mathbb{C}P^2\#8\overline{\mathbb{C}P}^2$ also admits infinitely many distinct smooth structures. We first construct a family of simply connected smooth 4-manifolds by modifying Stipsicz and Szabó's exotic 4-manifolds with $b_2^+ = 3$ and $b_2^- = 8$ in [22]. And then we compute their Seiberg-Witten invariants to show that they are mutually nondiffeomorphic.

2000 *Mathematics Subject Classification.* 57R17, 57R57, 57N13.

Key words and phrases. Exotic smooth structure, rational blow-down, Seiberg-Witten invariant.

* This work was supported by grant No. R01-2005-000-10625-0 from the KOSEF.

* The author also holds a joint appointment in the Research Institute of Mathematics, SNU.

Theorem 1.1. *The simply connected topological 4-manifold $3\mathbb{C}P^2 \# 8\overline{\mathbb{C}P}^2$ admits infinitely many distinct smooth structures.*

Acknowledgement. The author would like to thank András Stipsicz for his invaluable help and advice while working on this problem. He also pointed out critical errors in the early versions of this paper to the author.

2. A MAIN CONSTRUCTION

In this section we first briefly review a rational blow-down surgery and state related theorems ([3], [14] for details). Suppose that $p > q > 0$ are relatively prime integers. Let $C_{p,q}$ be a smooth 4-manifold obtained by plumbing disk bundles over the 2-sphere instructed by the following linear diagram

$$\begin{array}{ccccccc} -b_k & & -b_{k-1} & & \dots & & -b_1 \\ \bullet & \text{---} & \bullet & \text{---} & \dots & \text{---} & \bullet \\ u_k & & u_{k-1} & & & & u_1 \end{array}$$

where $\frac{p^2}{pq-1} = [b_k, b_{k-1}, \dots, b_1]$ is the unique continued fraction with all $b_i \geq 2$ and each vertex represents a disk bundle over the 2-sphere u_i whose Euler number is $-b_i$. Then $C_{p,q}$ is a negative definite simply connected 4-manifold whose boundary is a lens space $L(p^2, 1-pq)$. It was known that a lens space $L(p^2, 1-pq)$ bounds a rational ball $B_{p,q}$ with $\pi_1(B_{p,q}) \cong \mathbf{Z}_p$ ([1]). Furthermore, the inclusion $\partial B_{p,q} \rightarrow B_{p,q}$ induces an epimorphism $\pi_1(\partial B_{p,q}) \cong \mathbf{Z}_{p^2} \rightarrow \pi_1(B_{p,q}) \cong \mathbf{Z}_p$.

Definition Suppose X is a smooth 4-manifold containing a configuration $C_{p,q}$. Then we may construct a new smooth 4-manifold $X_{p,q}$, called the (*generalized*) *rational blow-down* of X , by replacing $C_{p,q}$ with the rational ball $B_{p,q}$. Note that this process is well-defined, that is, a new smooth 4-manifold $X_{p,q}$ is uniquely determined (up to diffeomorphism) from X because each diffeomorphism of $\partial B_{p,q}$ extends over the rational ball $B_{p,q}$. We call this a *rational blow-down surgery*.

A rational blow-down surgery with $q = 1$ was originally introduced by Fintushel and Stern to compute the Donaldson series for simply connected regular elliptic surfaces with multiple fibers of relatively prime orders ([3]) and later it was extended to the general case by the author ([14]). Furthermore, it was proved that the Seiberg-Witten basic classes and the Seiberg-Witten invariants of a smooth 4-manifold $X_{p,q}$ obtained by a rational blow-down surgery are closely related to those of X . For example, we have the following results.

Theorem 2.1 ([14]). *Suppose X is a smooth 4-manifold with $b_2^+ > 1$ which contains a configuration $C_{p,q}$. If L is a SW-basic class of X which satisfies $(L|_{C_{p,q}})^2 = -b_2(C_{p,q})$ and $c_1(L|_{L(p^2, 1-pq)}) = mp \in \mathbf{Z}_{p^2} \cong H^2(L(p^2, 1-pq); \mathbf{Z})$ with $m \equiv (p-1) \pmod{2}$, then L induces a SW-basic class $\overline{L}$ of $X_{p,q}$ such that $SW_{X_{p,q}}(\overline{L}) = SW_X(L)$.*

Corollary 2.1. *Suppose X is a smooth 4-manifold with $b_2^+ > 1$ which contains a configuration $C_{p,q}$. If L is a SW-basic class of X satisfying $L \cdot u_i = (b_i - 2), \forall 1 \leq i \leq k$ (or $L \cdot u_i = -(b_i - 2), \forall i$), then L induces a SW-basic class $\overline{L}$ of $X_{p,q}$ such that $SW_{X_{p,q}}(\overline{L}) = SW_X(L)$.*

Next, we review the exotic smooth 4-manifolds with $b_2^+ = 3$ and $b_2^- = 8$ constructed by Stipsicz and Szabó ([22] for details).

Let $E(2)$ be the simply connected elliptic surface with no multiple fibers and holomorphic Euler characteristic 2. Then $E(2)$ is an elliptic fibration over S^2 which has various types of singular fibers. In particular, it can be viewed as an elliptic fibration over S^2 with an I_{16} -singular fiber and eight fishtail-singular fibers ([8]). Note that one can change the eight fishtail fibers to three pairs of the same type fishtail fibers and two other fishtail fibers in $E(2)$. Let V_{K_1, K_2, K_3} be a smooth 4-manifold obtained by doing three knot surgeries in the three double node neighborhoods, each of which contains a pair of the same type fishtail fibers, in $E(2)$ respectively, using the twist knots K_1 for the first, K_2 for the second and K_3 for the third surgery. Then the smooth 4-manifold V_{K_1, K_2, K_3} contains a ‘pseudo-section’ i.e. an immersed 2-sphere with three positive double points and with square -2 ([5]). Furthermore, resolving the two positive intersections of this pseudo-section with the remaining two fishtail fibers, the 4-manifold V_{K_1, K_2, K_3} contains an immersed 2-sphere with five positive double points and with square 2. By blowing up at the five positive double points of the immersed 2-sphere, $V_{K_1, K_2, K_3} \# 5\overline{\mathbb{C}P}^2$ contains an embedded 2-sphere S of square -18 which still intersects the I_{16} -fiber transversally at one point. Let Σ_0 be an embedded 2-sphere in I_{16} -fiber which intersects S and let Σ_1 be an embedded 2-sphere in I_{16} -fiber which intersects with Σ_0 at one point, say q , positively. Then, applying 17 infinitely close blow-ups at the point q , the 4-manifold $V_{K_1, K_2, K_3} \# 22\overline{\mathbb{C}P}^2$ contains a configuration $C_{305, 17}$ which is a chain of embedded 2-spheres according to the following linear plumbing

$$(-18, -19, \overbrace{-2, \dots, -2}^{14}, -3, \overbrace{-2, \dots, -2}^{16}).$$

Note that the boundary $\partial C_{305, 17}$ is the lens space $L(305^2, -5184)$ which also bounds a rational ball $B_{305, 17}$. Hence, by rationally blowing down along the configuration $C_{305, 17}$ in $V_{K_1, K_2, K_3} \# 22\overline{\mathbb{C}P}^2$, Stipsicz and Szabó constructed a new family of smooth 4-manifolds

$$Y_{K_1, K_2, K_3} = (V_{K_1, K_2, K_3} \# 22\overline{\mathbb{C}P}^2 - \text{int } C_{305, 17}) \cup_{L(305^2, -5184)} B_{305, 17}.$$

Furthermore, they computed the Seiberg-Witten invariants of these manifolds. In particular, they showed that an infinite family of smooth 4-manifolds $Y_n := Y_{T_n, T_n, T_n}$ are pairwise nondiffeomorphic, all with nonvanishing Seiberg-Witten invariants.

Theorem 2.2 ([22]). *There are infinitely many pairwise nondiffeomorphic smooth, closed 4-manifolds Y_n with vanishing first homology, $b_2^+ = 3$, $b_2^- = 8$ and nontrivial Seiberg-Witten invariants.*

Now we will modify their construction to get an infinite family of simply connected smooth 4-manifolds with $b_2^+ = 3$ and $b_2^- = 8$. The procedure is the following: Recall that the 4-manifold V_{K_1, K_2, K_3} above contains a ‘pseudo-section’ i.e. an immersed 2-sphere with three positive double points and with square -2 . At this stage, instead of resolving two positive intersections of this pseudo-section with the remaining two fishtail fibers, resolve just one positive intersection point. Then the 4-manifold V_{K_1, K_2, K_3} contains an immersed 2-sphere with four positive double points and with square 0. By blowing up at the four positive double points of the immersed 2-sphere, $V_{K_1, K_2, K_3} \# 4\overline{\mathbb{C}P}^2$ contains an embedded 2-sphere of square -16 which still intersects one remaining fishtail fiber at one point, say p , and also intersects transversally the I_{16} -fiber. Now applying 2

infinitely close blow-ups at the point p and blowing up at the positive double point lying in the remaining fishtail fiber, we get a 4-manifold $V_{K_1, K_2, K_3} \# 7\overline{\mathbb{C}P}^2$ which contains an embedded 2-sphere S of square -18 and a configuration $C_{3,1}$ disjoint from S . And then, proceeding as above, i.e. applying 17 infinitely close blow-ups at the point q , we obtain a smooth 4-manifold $V_{K_1, K_2, K_3} \# 24\overline{\mathbb{C}P}^2$ which contains two disjoint configurations $C_{3,1}$ and $C_{305,17}$. More precisely, the 4-manifold $V_{K_1, K_2, K_3} \# 24\overline{\mathbb{C}P}^2$ contains a chain of embedded 2-spheres according to the following linear plumbing

$$(1) \quad (-5, -2, -1, -18, -19, \overbrace{-2, \dots, -2}^{14}, -3, \overbrace{-2, \dots, -2}^{16}).$$

Note that the third embedded 2-sphere, say E_6 , of square -1 in the linear plumbing above is an exceptional curve coming from the 6th blowing up of V_{K_1, K_2, K_3} . Since two configurations $C_{3,1}$ and $C_{305,17}$ are disjoint in $V_{K_1, K_2, K_3} \# 24\overline{\mathbb{C}P}^2$, we first get a family of smooth 4-manifolds, say Y'_{K_1, K_2, K_3} , by rationally blowing down along $C_{305,17}$, and then we finally get an infinite family of new smooth 4-manifolds, say Z_{K_1, K_2, K_3} , by rationally blowing down along $C_{3,1}$. Alternatively, we can get a family of smooth 4-manifolds, say Y''_{K_1, K_2, K_3} , by rationally blowing down along $C_{3,1}$ first, and then we can get the same smooth 4-manifolds Z_{K_1, K_2, K_3} by rationally blowing down along $C_{305,17}$. That is, we have

$$(2) \quad Y'_{K_1, K_2, K_3} = (V_{K_1, K_2, K_3} \# 24\overline{\mathbb{C}P}^2 - \text{int } C_{305,17}) \cup_{L(305^2, -5184)} B_{305,17}$$

$$(3) \quad Y''_{K_1, K_2, K_3} = B_{3,1} \cup_{L(9, -2)} (V_{K_1, K_2, K_3} \# 24\overline{\mathbb{C}P}^2 - \text{int } C_{3,1})$$

$$(4) \quad Z_{K_1, K_2, K_3} = B_{3,1} \cup_{L(9, -2)} (Y'_{K_1, K_2, K_3} - \text{int } C_{3,1})$$

$$(5) \quad = (Y''_{K_1, K_2, K_3} - \text{int } C_{305,17}) \cup_{L(305^2, -5184)} B_{305,17}.$$

Next, we will prove that the 4-manifolds Z_{K_1, K_2, K_3} constructed above are simply connected by using Van-Kampen's theorem.

Theorem 2.3. *For any twisted knots K_1, K_2 and K_3 , the 4-manifold Z_{K_1, K_2, K_3} is homeomorphic to $3\mathbb{C}P^2 \# 8\overline{\mathbb{C}P}^2$.*

Proof. First, in order to prove the simple connectivity of Z_{K_1, K_2, K_3} , let us decompose the smooth 4-manifold $V_{K_1, K_2, K_3} \# 24\overline{\mathbb{C}P}^2$ into $C_{3,1} \cup_{L(9, -2)} X_0 \cup_{L(305^2, -5184)} C_{305,17}$. Then the rational blow-down 4-manifolds Y'_{K_1, K_2, K_3} , Y''_{K_1, K_2, K_3} and Z_{K_1, K_2, K_3} can be decomposed as follows:

$$(6) \quad Y'_{K_1, K_2, K_3} = C_{3,1} \cup_{L(9, -2)} X_0 \cup_{L(305^2, -5184)} B_{305,17}$$

$$(7) \quad Y''_{K_1, K_2, K_3} = B_{3,1} \cup_{L(9, -2)} X_0 \cup_{L(305^2, -5184)} C_{305,17}$$

$$(8) \quad Z_{K_1, K_2, K_3} = B_{3,1} \cup_{L(9, -2)} X_0 \cup_{L(305^2, -5184)} B_{305,17}.$$

Let $i'_* : \pi_1(\partial C_{305,17}) \rightarrow \pi_1(C_{3,1} \cup_{L(9, -2)} X_0)$ and $i''_* : \pi_1(\partial C_{3,1}) \rightarrow \pi_1(X_0 \cup_{L(305^2, -5184)} C_{305,17})$ be induced homomorphisms by inclusions $i' : \partial C_{305,17} \rightarrow C_{3,1} \cup_{L(9, -2)} X_0$ and $i'' : \partial C_{3,1} \rightarrow X_0 \cup_{L(305^2, -5184)} C_{305,17}$ respectively. We may also choose the generators, say α and β , of $\pi_1(\partial C_{305,17}, x_0) \cong \mathbf{Z}_{305^2}$ and $\pi_1(\partial C_{3,1}, x_1) \cong \mathbf{Z}_9$ so that α and β are represented by circles lying in $\partial C_{305,17} \cap E_6$ and $\partial C_{3,1} \cap E_6$, respectively. Recall that E_6 is an exceptional curve which intersects transversally each embedded 2-sphere ending in the configurations $C_{3,1}$ and $C_{305,17}$ (refer to the linear plumbing (1) above). Since

$V_{K_1, K_2, K_3} \# 24\overline{\mathbb{C}P}^2$ is simply connected and since the images $i'_*(\alpha)$ and $i''_*(\beta)$ of generators are bounded by disks coming from a punctured exceptional curve $E_6 \setminus \{\text{an open disk}\}$, the two 4-manifolds Y'_{K_1, K_2, K_3} and Y''_{K_1, K_2, K_3} are simply connected. Hence, applying Van-Kampen's theorem on the decompositions (6) and (7) above, we conclude that both quotients groups

$$(9) \quad \pi_1(X_0 \cup_{L(305^2, -5184)} B_{305, 17}, x_1) / N_{j'_*(\beta)} \quad \text{and} \quad \pi_1(B_{3, 1} \cup_{L(9, -2)} X_0, x_0) / N_{j''_*(\alpha)}$$

are trivial. Here $j' : \partial C_{3, 1} \rightarrow X_0 \cup_{L(305^2, -5184)} B_{305, 17}$ and $j'' : \partial C_{305, 17} \rightarrow B_{3, 1} \cup_{L(9, -2)} X_0$ are inclusions, and $N_{j'_*(\beta)}$ and $N_{j''_*(\alpha)}$ are the least normal subgroups containing $j'_*(\beta)$ and $j''_*(\alpha)$ respectively. Note that they satisfy $j'_*(\beta)^9 = 1$ and $j''_*(\alpha)^{305^2} = 1$. Furthermore, there is a relation between $j'_*(\beta)$ and $j''_*(\alpha)$ when we restrict them to X_0 . That is, if we choose a path γ connecting x_0 and x_1 lying in $E_6 \setminus \{\text{two open disks}\}$, then they satisfy either $j'_*(\beta) = \gamma^{-1} \cdot j''_*(\alpha) \cdot \gamma$ or $j'_*(\beta) = \gamma^{-1} \cdot j''_*(\alpha)^{-1} \cdot \gamma$ (depending on orientations) because one is homotopic to the other in $E_6 \setminus \{\text{two open disks}\} \subset X_0$. Hence, by combining two facts above, we get $j'_*(\beta)^{305^2} = (\gamma^{-1} \cdot j''_*(\alpha)^{\pm 1} \cdot \gamma)^{305^2} = \gamma^{-1} \cdot j''_*(\alpha)^{\pm 305^2} \cdot \gamma = 1 = j'_*(\beta)^9$. Since the two numbers 9 and $305^2 = 25 \times 61^2$ are relatively prime, the element $j'_*(\beta)$ should be trivial. So the relation $j'_*(\beta) = \gamma^{-1} \cdot j''_*(\alpha)^{\pm 1} \cdot \gamma$ implies the triviality of $j''_*(\alpha)$. Finally the relation (9) above implies that both groups

$$(10) \quad \pi_1(X_0 \cup_{L(305^2, -5184)} B_{305, 17}) \quad \text{and} \quad \pi_1(B_{3, 1} \cup_{L(9, -2)} X_0)$$

are trivial. Thus, applying Van-Kampen's theorem again on the decomposition (8) above, we conclude that the 4-manifold Z_{K_1, K_2, K_3} is simply connected.

The rest of proof follows from simple Euler characteristics and signature computations together with Freedman's classification theorem. $\square$

Finally, we compute the Seiberg-Witten invariants of smooth 4-manifolds Z_{K_1, K_2, K_3} to show that they are mutually nondiffeomorphic. For simplicity, we restrict our attention to the special case $K_1 = K_2 = K_3 = n$ -twist knot T_n . Let V_n, Y'_n, Y''_n and Z_n denote simply connected smooth 4-manifolds $V_{T_n, T_n, T_n}, Y'_{T_n, T_n, T_n}, Y''_{T_n, T_n, T_n}$ and Z_{T_n, T_n, T_n} constructed above, respectively. Then the Seiberg-Witten invariants of Z_n can be easily computed using the same technique appeared in the proof of Theorem 3.3 in [22].

Theorem 2.4. *The smooth 4-manifold Z_n has only one (up to sign) SW-basic class $\overline{L}_n$ with $SW_{Z_n}(\overline{L}_n) = \pm n^3$.*

Proof. Since the Seiberg-Witten function of the $K3$ surface is $SW_{K3} = 1$ and the Alexander polynomial of T_n is $\Delta_{T_n} = nt - (2n - 1) + nt^{-1}$, by applying Theorem 1.1 in [4], the Seiberg-Witten function of the $V_n = V_{T_n, T_n, T_n}$ is equal to

$$SW_{V_n} = (ne^{2T} - (2n - 1) + ne^{-2T})^3,$$

where T is a regular elliptic fiber of V_n . And the blow-up formula implies that the Seiberg-Witten function of the $V_n \# 24\overline{\mathbb{C}P}^2$ is also equal to

$$SW_{V_n \# 24\overline{\mathbb{C}P}^2} = SW_{V_n} \cdot (e^{E_1} + e^{-E_1}) \cdots (e^{E_{24}} + e^{-E_{24}}),$$

where E_i is an exceptional curve coming from the i^{th} blowing up of V_n . Now applying Theorem 2.1 and Corollary 2.1 above for a rational blow-down 4-manifold Y'_n , we know that only the following SW-basic classes of $V_n \# 24\overline{\mathbb{C}P}^2$

$$L_n := \pm(6T + E_1 + \cdots + E_6 \pm E_7 + E_8 + \cdots + E_{24})$$

induce SW-basic classes L'_n of Y'_n and they all have SW-invariants $SW_{Y'_n}(L'_n) = \pm n^3$. Furthermore, applying Theorem 3.2 in [14] for Y'_n , we conclude that such L'_n 's are the only SW-basic classes of Y'_n . Next, again applying Theorem 2.1 and Corollary 2.1 above and Theorem 3.2 in [14] for the rational blow-down 4-manifold Z_n , we conclude that Z_n has only one (up to sign) SW-basic class $\bar{L}_n$ which is induced from the following SW-basic class of $V_n \# 24\overline{\mathbb{CP}}^2$

$$L_n = \pm(6T + E_1 + \cdots + E_6 + E_7 + E_8 + \cdots + E_{24})$$

and it has SW-invariant $SW_{Z_n}(\bar{L}_n) = \pm n^3$. $\square$

Corollary 2.2. *The 4-manifolds Z_n ($n \geq 1$) are simply connected irreducible smooth 4-manifolds which are mutually nondiffeomorphic.*

Proof of Theorem 1.1. By Theorem 2.3 and Corollary 2.2 above, the 4-manifolds Z_n with $n \geq 1$ provide an infinite family of smooth 4-manifolds which are homeomorphic, but not diffeomorphic, to $3\mathbb{CP}^2 \# 8\overline{\mathbb{CP}}^2$. Hence we are done. $\square$

3. MORE EXAMPLES

In this section we construct more examples of simply connected smooth 4-manifolds with $b_2^+ = 3$ and $b_2^- = 8$ which are mutually nondiffeomorphic by using a different configuration.

First recall that the 4-manifold V_{K_1, K_2, K_3} contains a ‘pseudo-section’ with three positive double points and with square -2 . At this time, by blowing up at the three positive double points of the pseudo-section, we get a 4-manifold $V_{K_1, K_2, K_3} \# 3\overline{\mathbb{CP}}^2$ which contains an embedded 2-sphere of square -14 and which intersects two remaining fishtail fibers at the points p_1 and p_2 respectively. Applying 2 infinitely close blow-ups at p_1 and p_2 respectively and blowing up at two positive double points lying in the two remaining fish-tail fibers, we get a 4-manifold $V_{K_1, K_2, K_3} \# 9\overline{\mathbb{CP}}^2$ which contains an embedded 2-sphere S of square -18 and two copies, say $C_{3,1}$ and $C'_{3,1}$, of a configuration $C_{3,1}$ disjoint from S . And then, applying 17 infinitely close blow-ups at the point q , we obtain a smooth 4-manifold $V_{K_1, K_2, K_3} \# 26\overline{\mathbb{CP}}^2$ which contains three disjoint configurations $C_{3,1}$, $C'_{3,1}$ and $C_{305,17}$. Since three configurations are disjoint in $V_{K_1, K_2, K_3} \# 26\overline{\mathbb{CP}}^2$, we get a new family of smooth 4-manifolds $\tilde{Y}_{K_1, K_2, K_3}$, $\tilde{Y}'_{K_1, K_2, K_3}$ and $\tilde{Z}_{K_1, K_2, K_3}$ by rationally blowing down along $C_{3,1}$, $C'_{3,1}$ and $C_{305,17}$ sequentially. Alternatively, we can get a family of smooth 4-manifolds $\tilde{Y}''_{K_1, K_2, K_3}$, $\tilde{Y}'''_{K_1, K_2, K_3}$ and $\tilde{Z}_{K_1, K_2, K_3}$ by rationally blowing down along $C_{305,17}$, $C_{3,1}$ and $C'_{3,1}$ sequentially. Then, using the same techniques as in the proof of Theorem 2.3 and Theorem 2.4 in Section 2, we can prove that

Theorem 3.1. *For any twisted knots K_1, K_2 and K_3 , the 4-manifold $\tilde{Z}_{K_1, K_2, K_3}$ is homeomorphic to $3\mathbb{CP}^2 \# 8\overline{\mathbb{CP}}^2$.*

Theorem 3.2. *The smooth 4-manifold $\tilde{Z}_n = \tilde{Z}_{T_n, T_n, T_n}$ has only one (up to sign) SW-basic class $\bar{L}_n$ with $SW_{\tilde{Z}_n}(\bar{L}_n) = \pm n^3$.*

Corollary 3.1. *The 4-manifolds $\tilde{Z}_n$ ($n \geq 1$) are simply connected irreducible smooth 4-manifolds with $b_2^+ = 3$ and $b_2^- = 8$ which are mutually nondiffeomorphic.*

Remark. It is an interesting question to ask whether Z_{K_1, K_2, K_3} and $\tilde{Z}_{K_1, K_2, K_3}$ constructed in this paper are diffeomorphic to the Stipsicz and Szabó's exotic 4-manifolds with $b_2^+ = 3$ and $b_2^- = 8$ in [22]. Note that we still do not know whether the Stipsicz and Szabó's exotic 4-manifolds are simply connected or not.

REFERENCES

- [1] A. Casson and J. Harer, *Some homology lens spaces which bound rational homology balls*, Pacific. J. Math. **96** (1981), 23–36
- [2] S. Donaldson, *Irrationality and the h-cobordism conjecture*, J. Diff. Geom. **26** (1987), 141–168
- [3] R. Fintushel and R. Stern, *Rational blowdowns of smooth 4-manifolds*, J. Diff. Geom. **46** (1997), 181–235
- [4] R. Fintushel and R. Stern, *Knots, links and 4-manifolds*, Invent. Math. **134** (1998), 363–400
- [5] R. Fintushel and R. Stern, *Double node neighborhoods and families of simply connected 4-manifolds with $b^+ = 1$* , to appear in J. Amer. Math. Soc. (arXiv:math.GT/0412126)
- [6] R. Friedman and J. Morgan, *Smooth 4-manifolds and complex surfaces*, Ergebnisseder Mathematik und ihrer Grenzgebiete **27**, Springer-Verlag, 1994
- [7] R. Gompf, *A new construction of symplectic manifolds*, Annals of Math. **142** (1995), 527–595
- [8] J. Harer, A. Kas and R. Kirby, *Handlebody decompositions of complex surfaces*, Memoirs of Amer. Math. Soc. **62**, 1986
- [9] D. Kotschick, *On manifolds homeomorphic to $\mathbb{C}P^2\#8\overline{\mathbb{C}P}^2$* , Invent. Math. **95** (1989), 591–600
- [10] J. Morgan and K. O'Grady, *Differential topology of complex surfaces. Elliptic surfaces with $p_g = 1$: smooth classification*, Lecture Notes in Mathematics **1545**, Springer-Verlag, 1993
- [11] D. Park, *Exotic smooth structures on $\mathbb{C}P^2\#n\overline{\mathbb{C}P}^2$* , Proc. Amer. math. Soc. **128** (2000), 3057–3065
- [12] D. Park, *Exotic smooth structures on $\mathbb{C}P^2\#n\overline{\mathbb{C}P}^2$, Part II*, Proc. Amer. math. Soc. **128** (2000), 3067–3073
- [13] D. Park, *Constructing infinitely many smooth structures on $3\mathbb{C}P^2\#n\overline{\mathbb{C}P}^2$* , Math. Ann. **322** (2000), 267–278
- [14] J. Park, *Seiberg–Witten invariants of generalized rational blow-downs*, Bull. Austral. Math. Soc. **56** (1997), 363–384
- [15] J. Park, *Exotic smooth structures on 4-manifolds*, Forum Math. **14** (2002), 915–929
- [16] J. Park, *Exotic smooth structures on 4-manifolds, II*, Topology Appl. **132** (2003), 195–202
- [17] J. Park, *Simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$* , Invent. Math. **159** (2005), 657–667
- [18] J. Park, A. Stipsicz and Z. Szabó, *Exotic smooth structures on $\mathbb{C}P^2\#5\overline{\mathbb{C}P}^2$* , Math. Res. Letters **12** (2005), 701–712
- [19] A. Stipsicz, *Donaldson series and (-1) -tori*, J. Reine Angew. Math. **465** (1995), 133–144
- [20] A. Stipsicz and Z. Szabó, *The smooth classification of elliptic surfaces with $b_2^+ > 1$* , Duke Math. J. **75** (1994), 1–50
- [21] A. Stipsicz and Z. Szabó, *An exotic smooth structure on $\mathbb{C}P^2\#6\overline{\mathbb{C}P}^2$* , Geometry & Topology **9** (2005), 813–832
- [22] A. Stipsicz and Z. Szabó, *Small exotic 4-manifolds with $b_2^+ = 3$* , arXiv:math.GT/0501273
- [23] Z. Szabó, *Irreducible four-manifolds with small Euler characteristics*, Topology **35** (1996), 411–426

DEPARTMENT OF MATHEMATICAL SCIENCES, SEOUL NATIONAL UNIVERSITY, SAN 56-1 SILLIM-DONG, GWANAK-GU, SEOUL 151-747, KOREA

E-mail address: jipark@math.snu.ac.kr